

This item is the archived peer-reviewed author-version of:

Enhancement of co-production of nutritional protein and carotenoids in *Dunaliella salina* using a two-phase cultivation assisted by nitrogen level and light intensity

Reference:

Sui Yixing, Muys Maarten, Van de Waal Dedmer, D' Adamo Sarah, Vermeir Pieter, Fernandes Tânia V., Vlaeminck Siegfried.- Enhancement of co-production of nutritional protein and carotenoids in *Dunaliella salina* using a two-phase cultivation assisted by nitrogen level and light intensity

Bioresource technology - ISSN 0960-8524 - 287(2019), 121398

Full text (Publisher's DOI): <https://doi.org/10.1016/J.BIORTECH.2019.121398>

To cite this reference: <https://hdl.handle.net/10067/1596610151162165141>

Enhancement of co-production of nutritional protein and carotenoids in *Dunaliella salina* using a two-phase cultivation assisted by nitrogen level and light intensity

Yixing Sui¹, Maarten Muys¹, Dedmer B. Van de Waal², Sarah D'Adamo³, Pieter Vermeir⁴,
Tânia V. Fernandes² and Siegfried E. Vlaeminck^{1*}

¹Research Group of Sustainable Energy, Air and Water Technology, Department of Bioscience Engineering, University of Antwerp, Groenenborgerlaan 171, 2020 Antwerpen, Belgium

²Department of Aquatic Ecology, Netherlands Institute of Ecology (NIOO-KNAW), PO Box 50, 6700 AB Wageningen, The Netherlands

³Bioprocess Engineering, Wageningen University & Research, PO Box 16, 6700 AA, Wageningen, The Netherlands

⁴Laboratory of Chemical Analysis, Department of Green Chemistry and Technology, Ghent University, Valentin Vaerwyckweg 1, 9000 Gent, Belgium

*: Corresponding author: siegfried.vlaeminck@uantwerpen.be

Abstract

Microalga *Dunaliella salina* is known for its carotenogenesis. At the same time, it can also produce high-quality protein. The optimal conditions for *D. salina* to co-produce intracellular pools of both compounds, however, are yet unknown. This study investigated a two-phase cultivation strategy to optimize combined high-quality protein and carotenoid production of *D. salina*. In phase-one, a gradient of nitrogen concentrations was tested. In phase-two, effects of nitrogen pulse and high illumination were tested. Results reveal optimized protein quantity, quality (expressed as essential amino acid index EAAI) and carotenoids content in a two-phase cultivation, where short nitrogen starvation in phase-one was followed by high illumination during phase-two. Adopting this strategy, productivities of protein, EAA and carotenoids reached 22, 7 and 3 mg/L/d, respectively, with an EAAI of 1.1. The quality of this biomass surpasses FAO/WHO standard for human nutrition, and the observed level of β -carotene presents high antioxidant pro-vitamin A activity.

Key words

Single-cell protein; pigment; nitrogen limitation; food; microalgae

1. Introduction

The global population will reach 9.3 billion by 2050, with 6.4 billion of people in urban areas (Corcoran et al., 2010). The societal changes of both population and living standard are leading to 50% increase of protein demand, and even 82% and 102% increase of dairy and meat products by 2050, respectively (Boland et al., 2013). Along with protein shortage, the deficiency of functional nutrients in food, like β -carotene, are causing severe health problems for human. Specifically, β -carotene is essential for the human body due to its antioxidant pro-vitamin A activity, and insufficient uptake of β -carotene will lead to severe vitamin A deficiency, prompting human blindness and affecting immune response systems (Sommer, 2001). Currently, β -carotene and vitamin A deficiency have become a major public health concern in more than 70 countries (Sommer, 2001). To sustainably fulfill the protein gap for human consumption, novel protein sources such as microalgae are considered important contributions (Muys et al., 2019). Moreover, microalgae with elevated carotenogenesis can further increase nutritional quality by preventing vitamin A deficiency. Particularly, natural β -carotene found in microalgae, fruits and vegetables has the advantage of its mixed stereoisomers of *all-trans* and *9-cis* β -carotene, which are more fat-soluble and less crystallizable than synthetic β -carotene (*all-trans* β -carotene) (Ben-Amotz, 1993). By consuming carotenoid-rich diet, potentially lower incidence of various kinds of cancer can be expected (Ben-Amotz, 1993)

Microalgal production is conventionally aiming at one specific target, such as biomass from *Chlorella*, protein from *Spirulina* and β -carotene from *Dunaliella* (Ben-Amotz, 1993).

Consequently, individual production lines are required to achieve production of multiple target compounds. If one microalgal species possesses the ability to optimally co-produce both high-quality protein as well as β -carotene at the same time, the efficiency in production of multiple high-value products can be substantially increased.

It is well established that the microalga *Dunaliella salina* is one of the best sources of natural β -carotene, which can contribute up to 14% of the cell dry weight (Aasen et al., 1969). Up to date, many studies have shown that stress conditions are major factors enhancing the accumulation of β -carotene in *D. salina*, such as high light intensity, high salinity, extreme temperatures and nitrogen (N) deficiency, as a protective mechanism to prevent cellular damages e.g. photo-damage (Lamers et al., 2008; Marín et al., 1998).

While carotenogenesis by *D. salina* has been widely studied, its potential as protein source has drawn limited attention. In fact, *D. salina* can display high protein content (up to 80% of ash free dry weight), which is subjected to different cultivation conditions and growth phases (Sui and Vlaeminck, 2019). Furthermore, the essential amino acid (EAA) content of *D. salina* fulfills human requirement as indicated by FAO/WHO reference, which defines its high-quality protein profile (Becker, 2007; Sui et al., 2019). When comparing with other microalgae such as *Chlorella* and *Spirulina*, both protein and EAA content of *D. salina* have either comparable or even superior values (Becker, 2007; Muys et al., 2019; Sui et al., 2019). Based on the characteristics of *D. salina*, it strongly appears to be a valuable candidate as novel food source, containing both high-quality protein and β -carotene.

Based on both lab- and large-scale experience with cultivating *Dunaliella* for β -carotene production, a two-phase cultivation system was proven to be successful, and has been applied by commercial producers (Ben-Amotz, 1995). The concept of such two-phase systems is to increase microalgal biomass level at phase-one with optimum growth conditions, and to induce β -carotene production at phase-two with enhanced stress conditions, e.g. high light intensity and N deficiency (Ben-Amotz, 1995). Moreover, it has been reported that stress conditions can also contribute to the up-regulation of EAA levels in plants and microalgae (Galili et al., 2016; Obata and Fernie, 2012). Specifically, high light intensity and N deficiency have been shown to enhance the production of EAA, especially lysine, threonine, methionine, valine and isoleucine in microalgae (Kiyota et al., 2014; Zhang et al., 2016). For *D. salina*, there is no report on the regulation of EAA affected by cultivation conditions, and the combined production of two main nutritional compounds from *D. salina* has not been studied yet (Sui et al., 2019). Instead of a process targeting either β -carotene or protein production, a two-phase cultivation approach adopting sequential stress conditions may prove to be an effective way to boost both high-quality protein and β -carotene production from *D. salina*.

In this study, the impact of a gradient in N availability together with N pulses and high light intensities on the dynamics of biomass, protein, EAA and β -carotene production in *D. salina* have been explored over different growth phases in a two-phase cultivation approach. This study intends to demonstrate for the first time an optimized cultivation

condition and harvest regime for the maximum production of both EAA and β -carotene of *D. salina*.

2. Materials and methods

2.1 Microalgal strain, two-phase cultivation approach and cultivation conditions

D. salina CCAP 19/18 was purchased from Culture Collection of Algae and Protozoa (CCAP, Scotland, UK). Sterilized Modified Johnson's medium (Borowitzka, 1988) as standard medium for *D. salina* with different N levels was used for cultivation. Six treatments covering a gradient of N concentrations were tested in phase-one (N1 to N6; Table 1). When the algae reached stationary phase, each treatment was divided into two further conditions for phase-two: high N (NN1 to NN6) and high light intensity (NL1 to NL6; Table 1). All treatments were performed in triplicates for both phases. Triplicates in phase-two were derived from the pooled triplicates of corresponding treatments in phase-one. The initial biomass concentration in phase-one was around 40,000 cells/mL. Experiments were conducted in a water bath with a controlled temperature of 25°C. Continuous light was provided by fluorescent tubes at an incident irradiance of 70 $\mu\text{mol photons/m}^2/\text{s}$ for standard conditions, and 110 $\mu\text{mol photons/m}^2/\text{s}$ for high irradiance conditions (Philips TL-D 30W/33-640, the Netherlands). Mixing and aeration were given by a mixture of pre-humidified air and 2% CO_2 . Although pH was not controlled, it was found stable (7.8-8) over the experiment due to the CO_2 addition (Table 1).

2.2 Sample analyses, calculations and statistics

Daily samples from all treatments were analyzed directly for cell number and cell volume, and preserved at -20°C for protein, carotenoids and EAA analyses at the end of the experiment. A Multisizer 3 Coulter Counter was used for both cell number and volume measurement. The protein content was determined following Markwell method, a modified Lowry method with bovine serum albumin as standard (Markwell et al., 1978). Total carotenoids content was measured according to Lichtenthaler, (1987) with 100% acetone extraction:

$$\text{Chlorophyll } a \text{ (mg/L)} = 11.24 \times OD_{661.6} - 2.04 \times OD_{644.8}$$

$$\text{Chlorophyll } b \text{ (mg/L)} = 20.13 \times OD_{644.8} - 4.19 \times OD_{661.6}$$

Total carotenoids (mg/L)

$$= \frac{1000 \times OD_{470} - 1.90 \times \text{Chlorophyll } a - 63.14 \times \text{Chlorophyll } b}{214}$$

where $OD_{661.6}$, $OD_{644.8}$ and OD_{470} refer to the optical densities of the extracted supernatant measured at 661.6 nm, 644.8 nm and 470 nm, respectively.

To prepare for EAA analysis, samples were centrifuged (5,000 x g, 10 min), hydrolyzed (6M HCl, 110°C, 24 hours) with vacuum and evaporated, after which samples were re-dissolving in 0.75mM HCl and stored at -20°C before analysis. For EAA determination, the EZ:faast amino acids analysis procedure was adopted (Phenomenex, 2003), with separation using gas chromatography (Agilent HP 6890, USA) and detection using mass spectrometry (Agilent HP 5973, USA). The essential amino acids index (EAAI) was derived

based on the EAA content with FAO/WHO EAA requirements for human as reference

(Oser, 1959; WHO/FAO/UNU Expert Consultation, 2007):

$$EAAI = \sqrt[n]{\frac{aa1}{AA1} \times \frac{aa2}{AA2} \times \dots \times \frac{aan}{AAn}}$$

where *aan* and *AAn* are the EAA content over total protein content (mg EAA/g protein) in the sample and FAO/WHO reference, respectively. An EAAI value of ≥ 1 , 0.95-1, 0.86-0.95, 0.75-0.86 and ≤ 0.75 indicates its superior quality, high quality, good quality, useful quality and inadequate quality, respectively (Kent et al., 2015). For comparison purposes, it has been calculated that the EAAI value of egg and soybean are 1.65 and 1.34 separately (Becker, 2007).

Total protein and carotenoids per liter of culture (mg/L) were defined as suspension protein and carotenoids content. The protein, carotenoid and EAA productivity (mg/L/d) were calculated from their suspension content (mg/L) divided by the time of cultivation (days) at each sampling point.

Samples at the end of the two phases were analyzed for nitrate (NO_3^-), ammonium (NH_4^+) and phosphate (PO_4^{3-}) concentrations in the medium. As ammonium concentrations were not detected in any treatment, they were not reported in this study. Filtered samples were diluted with de-ionized water accordingly and a Seal QUATRO Auto Analyzer (Seal Analytical Inc., the Netherlands) was used for determination following standard methods (APHA, 2012).

Multiple regression analysis in SPSS statistics 24 was used to compare data in Fig. 3. A significance level $p < 0.05$ was considered as statistically different. All results were expressed as means \pm standard deviations in tables and figures (apart from EAA and EAA derived parameters). The values stated in the main text were without standard deviation for better readability.

3. Results and discussion

3.1 Biomass growth

Both the N level and light intensity greatly affected the microalgal growth. During phase-one, the microalgal cells reached different concentrations at stationary growth phase from N1 to N4, ranging from approximately 2×10^5 cells/mL to 2.5×10^6 cells/mL. This was mostly contributed by the differences in initial N concentrations in the medium, which were depleted at the end of phase-one (Table 2). From N4 to N6, the cell densities did not change, despite different initial N concentrations (Fig 1A). Considering that there was still N remaining in the medium after phase-one for N5 and N6, the cells in these treatments were limited by another factor, while for N4 cells may have been co-limited. One likely limiting factor can be phosphorus (P), as the residual P in the medium from N4 to N6 was zero (Table 2). Besides, light limitation could occur as well, which will be discussed in more detail later.

During phase-two, both higher N and higher light intensity had effect on biomass growth. Specifically, from NN1 to NN3, cell numbers increased with extra N addition as a result

from their N starvation in phase-one. Differently, cell concentration from NN4 to NN6 did not increase, and even slightly decreased after N addition, indicating that N was not the limiting factor. Higher light intensity did not affect the cell concentration in NL1, most likely due to the N scarcity reached during phase-one. For NL2 and NL3, higher light intensity promoted cell growth, although N or P was limiting. It is possible that *D. saline* in NL2 and NL3 might have stored both N and P inside the cell to be used for further growth under higher light intensity (Dortch et al., 1984). From NL4 to NL6, higher light intensity did not affect much or slightly lowered the biomass, which demonstrates that light is not the limiting factor. At this point, P concentration in the medium is expected to be the main limiting factor which prohibited the further growth of cells (Table 2). This finding is in line with previous work, where higher light intensity could not further boost cell densities of *Arthrospira platensis* when P was depleted (Markou et al., 2012).

Besides cell densities, also cell sizes were subject to changes in response to the N and NN treatments (Fig. 1). An oscillation pattern was visible, indicating the variations of cell sizes at different growth stages. This pattern might be related to cell division, where cells grew exponentially in the exponential growth phase, and reached their maximum in stationary phase. These findings suggest that microalgal cells start increasing in size until they reach a critical point for cell division, which can be likely affected by external energy supply e.g. light, temperature and nutrients (Zachleder et al., 2016). This explains the less pronounced oscillation in NN5 and NN6 even when extra N was supplied, indicating again that cells were P limited. Nonetheless in this study, higher light intensity does not seem to

result in distinct cell volume changes (Fig 1A). It has been shown that cell oscillation by lighting can be species-specific, where cell volume changes are not restricted to a fixed pattern (Agusti and Kalff, 1989).

3.2 Protein and carotenoids dynamics

During phase-one, all treatments from N1 to N6 showed increases of suspension protein along with cell growth, and the more initial N in the medium, the more suspension protein was reached at the end of phase-one, ranging from 17 mg/L in N1 to 597 mg/L in N6 (Fig 1B). The corresponding cellular protein content also reached the highest level in N6 (233 pg/cell) from N1 (83 pg/cell). Higher protein levels at increased N concentrations in suspension, cell, and biomass have all been reported for various species such as *Dunaliella tertiolecta*, *Scenedesmus* sp. LX1, and *Chlorella* sp. (Fabregas et al., 1989; Kiran et al., 2016; Zhuang et al., 2018), while N starvation is well known to reduce the protein content (Gao et al., 2018).

During the high-N treatments in phase-two, both suspension protein and cellular protein were significantly boosted from NN1 to NN4. Such rises were observed very shortly after N addition and continued until the end of phase-two (Fig 1B). The biggest increase of suspension protein occurred in NN1 with 3460% (from 17 mg/L to 599 mg/L), which was due to the extremely low protein and biomass concentration in N1. The highest suspension protein at the end of phase-two reached 902 mg/L in NN4. For cellular protein, similar results were found with 404% increase in NN1 reaching 419 pg/cell, which is comparable with NN2-NN6. When microalgae are supplied with excess of substrate after a

period of starvation, overcompensating mechanisms can occur, in which cells are triggered to uptake and store higher substrate amounts than necessary (Brown and Shilton, 2014). The boost of protein production in NN1-NN4 can likely be a result of these mechanisms. Another event that can occur when enough N is present in the culture is called “luxury uptake”, which implies the natural uptake of resources beyond necessity without prior starvation. The increase in protein content of NN5-NN6 could be a result of this mechanism (Brown and Shilton, 2014). Both mechanisms linking to the survival of microalgae are commonly found and thoroughly described (Xie et al., 2017). Due to these two phenomena, protein or N content of many microalgal species such as *Chlorella vulgaris*, and also macroalgal species such as *Oedogonium* could be boosted with N addition (Cole et al., 2015; Dortch et al., 1982; Xie et al., 2017).

During the higher-light treatments in phase-two, however, both suspension and cellular protein level for NL1-NL4 did not substantially change, likely due to N depletion from the previous phase. In NL5 and NL6, however, where N was still present, suspension and cellular protein levels increased by up to 37% and 77%, respectively. Higher protein content induced by higher light intensity has been observed for *Chlorella vulgaris* and *Chlorella pyrenoidosa* (Chen et al., 2015; Ogbonna and Tanaka, 1996). However, other microalgal species could also present either decreased protein content or no change at all following higher light intensity, especially with low N availability (Chen et al., 2015; Markou et al., 2012). Thus, the results from both high-N and higher-light treatments suggest that N levels directly links to the total protein accumulation.

The dynamics of carotenoids generally showed the opposite pattern of protein. During phase-one from N1 to N4, both suspension and cellular carotenoids accumulated, with maximum 29 mg/L and 16 pg/cell reached during the stationary phases of N3 and N1, respectively (Fig 1C). This trend is a consequence of N deficiency, which is one of the most effective ways to induce β -carotene accumulation in *D. salina* (Lamers et al., 2012). The cellular carotenoids showed no changes in N5 and N6, although a slight increase of suspension carotenoids occurred, which was mainly due to an increase in biomass (Fig 1C). During the higher-N treatment in phase-two, all cellular carotenoids from NN1 to NN6 dropped to initial levels, around 4-8 pg/cell. Due to different biomass levels in this phase, the suspension carotenoids either dropped or slightly increased, but remained in a similar range from 9 to 22 pg/cell for all treatments (Fig 1C). During the higher-light treatment in phase-two, both cellular and suspension carotenoids increased rapidly in NL1 to NL4. The increases started shortly after the switch to higher light and continued till the end of the experiment, with maximum 196% cellular carotenoids increase in NL4 and 262% suspension carotenoids increase in NL2 (Fig 1C). Very little changes of carotenoids levels were found in NN5, NL5, NN6 and NL6, primarily due to the presence of N, which suggest N limitation to be the determining factor for effective carotenoids induction. When light intensity is higher than needed for photosynthesis in *D. salina*, β -carotene is produced in excess to overcome light stress and potential photo-oxidative damage, and it has been well documented that high light intensity can significantly increase the production of β -carotene in *D. salina* (Lamers et al., 2010; Raja et al., 2007). This is also one important

reason why large-scale cultivation of *D. salina* is located wherever high light intensity is expected (Ben-Amotz, 1993).

3.3 Protein and carotenoids productivities

In phase-one, protein productivities generally declined (N1 and N3), or rose and declined (N2, N4-N6) towards the stationary phase (Fig 1D). These results showed that higher protein productivity was obtained with high N availability. The highest protein productivities reached were 7, 28, 35, 50, 50 and 49 mg/L/d, for N1 to N6, respectively, and mostly occurred between exponential and linear growth phase. This is in agreement with earlier work on *D. salina* tested at different salinities, pH levels and light regimes (Sui et al., 2019; Sui and Vlaeminck, 2019). For treatments with N starvation at the end of phase-one, namely N1-N4, higher-N treatment in phase-two (NN1-NN4) significantly stimulated their protein productivity by 37-1024%. This again can be linked to N overcompensating mechanism. Differently, for N5 and N6, where N was still abundant after phase-one, protein productivity in phase-two (NN5-NN6) did not show notable trend changes but kept declining (Fig 1D). During the higher-light treatment in phase-two, all protein productivities continued to decrease, as a result from prolonged cultivation time and rather stable protein content (Fig 1B, 1D). Overall, these results show that when *D. salina* cells experience N starvation, extra N addition enhances protein productivity, likely due to overcompensating mechanisms. These findings can be used to optimize the design for harvesting strategy, which could increase biomass protein content. When abundant N is provided, the optimal harvesting point will be around exponential to linear growth

phase. This recommendation, however, only applies to maximize protein quantity without considering its quality.

Differently from protein productivity, carotenoids productivities showed consistent increase pattern during phase-one from N1 to N4, with a maximum of 3 mg/L/d reached in N3 (Fig 1E). These high carotenoids productivities followed the pattern of their suspension carotenoids contents, which depends on both biomass and cellular carotenoids accumulation (Fig 1C). During higher-light treatment in phase-two, NL1 to NL4 all showed significant increases of carotenoids productivity, ranging from 15 to 107%. The highest carotenoids productivity reached was 4 mg/L/d in NL3 (Fig 1E). In contrast, the high-N treatments in phase-two contributed negatively to all carotenoids productivities from NN1 to NN4, which decreased by 5 to 54% (Fig 1E). For the treatments of N5, NN5 and NL5, as well as N6, NN6 and NL6, low productivities without evident changes in carotenoids were observed throughout the experiment, which was possibly due to the presence of excess N. Generally, N starvation in *D. salina* enhances the accumulation of carotenoids, which further increases with higher-light exposure. Thus, the combination of N starvation and subsequent exposure to higher light can be a beneficial way to boost carotenoids production.

3.4 EAA dynamics and productivities

A few samples have been selected for a more detailed analyses of EAA. They were the intermediate treatment on the intersection of N and P limitation (N4) and the highest N treatment (N6) as reference, together with the respective high nitrogen (NN4 and NN6)

and light (NL4 and NL6) treatments, both just after the transfer (start) and at the end of the experiment (end). At the end of phase-one, N4 clearly showed higher levels of EAA content, EAA productivity and EAAI relative to N6 (Fig 2A). The EAAI of N4 reached 1.3, which is of superior quality for human consumption (threshold EAAI = 1), while the EAAI of N6 was only 0.4, showing inadequate quality. Looking at individual EAA levels, all the EAA in N4 were substantially present in higher amounts as compared to N6, exceeding FAO/WHO reference except for valine (Fig 2B, 2C and Table 3). These findings indicate that *D. salina* with EAAI of 1.3 is well suited to be incorporated into food which perfectly match human requirement (EAAI = 1), and actually saving 23% of biomass, further increasing the efficiency of food consumption. Protein and amino acid synthesis in microalgae naturally relies on N assimilation pathways, where nitrate is transported inside the cell and converted to nitrite by nitrate reductase. Ammonia is then obtained by further reduction of nitrite and incorporated into glutamate/glutamine via glutamine synthase and NADPH-dependent glutamine:2-oxoglutarate aminotransferase (GS/GOGAT) pathway (Alipanah et al., 2015; Halsey et al., 2011; Remmers et al., 2018; Sanz-Luque et al., 2015). Glutamate/glutamine sequentially provides the critical entry point of N into cellular biochemicals, which can subsequently be used for synthesis of other EAAs (Guerra et al., 2013). Although N availability associates closely with EAA production in microalgae, as shown in this study and many others, a higher N level does not necessarily lead to higher EAA production. When microalgal cells become N limited, mostly towards the stationary growth phase, their major response is to preserve cellular N capacity via scavenging

mechanisms, by which EAA biosynthesis can still occur using intracellular N (Alipanah et al., 2015; Halsey et al., 2011; Lv et al., 2017; Remmers et al., 2018; Zhang et al., 2016).

However, when N deprivation occurs in the long term, EAA synthesis is interrupted and results in sharp decreases of protein and amino acids (Kiyota et al., 2014; Van de Waal et al., 2010; Zhang et al., 2016).

Several studies have shown that N limitation or a short period of starvation can boost EAA production in microalgae and macroalgae. For instance, it has been shown that all EAA levels of *D. salina* SAG 184.80 were enhanced towards stationary growth phase, when N became limited and then shortly starved (Sui et al., 2019). *Synechocystis* sp. also exhibited rising levels of all EAA during short N starvation, while longer N starvation resulted in dropping of EAA levels (Kiyota et al., 2014). Similarly, short N starvation contributed to the production of several EAAs, especially phenylalanine, in marine microalga *Isochrysis zhangjiangensis*, which can be the consequence of N scavenging of e.g. nucleotides and rubisco protein. Nevertheless, significant decreased EAA levels were found after long-exposed N deficiency (Zhang et al., 2016). Besides, the macroalgae *Ulva ohnoi* was also shown to exhibit higher proportions of alanine, serine, glycine, and the EAAs phenylalanine, threonine and valine with low N concentration (Angell et al., 2014).

When extra N was supplied during phase-two, EAA content, EAA productivity and EAAI from NN4 (start) decreased sharply, resulting in inadequate protein quality (Fig 2A). A similar drop was also observed in NN6 (start). Regarding individual EAA levels, higher N addition also led to their reductions, where the overall level of EAA decreased

dramatically from N4 to NN4 (start and end) (Fig 2B and Table 3). Overall, these results clearly demonstrate that short N starvation promotes EAA accumulation in *D. salina*, while high N levels has a negative impact. At the end of phase-two after N addition, EAA content, EAA productivity, and EAAI in NN4 (end) and NN5 (end) all increased by 14-28%, 3-14%, and 35-55%, respectively. This again indicates that cells tend to preserve EAAs towards later growth phases to maintain the N capacity and cell functions, possibly due to the drop of N concentration in the environment (Fig 2A, Table 2).

While the complex biosynthesis of EAAs following various pathways are associated with N availability, it is not always easy to link them to other environmental conditions. Here, higher light intensity in phase-two seemed to have differential effects on EAA production. In the higher light treatment, NL4 (start and end) did not respond positively, resulting in slight decreases of EAA content, EAA productivity, and EAAI (Fig 2A). Nonetheless, protein quality of the biomass at this stage still remained superior for human consumption, as indicated by EAAI and relatively high amounts of individual EAAs (Fig 2A, 2B and Table 3). Although *D. salina* biomass seems to maintain protein quality during the higher-light treatment, thus only slight differences between NL4 (start) and NL4 (end) in EAA content, the EAA productivity in NL4 (end) decreased by 46%, which was mainly attributed to growth stage (Fig 2A). In NL6 (start and end), higher light intensity evidently promoted the accumulation of EAA in the biomass with higher EAA content (Fig 2A). Shortly after introducing higher light in NL6 (start), EAA content, EAA productivity and EAAI increased with 49, 25, and 30%, respectively. At the end of phase-two in NL6 (end), such increases

were even 232, 152, and 189%, reaching 36%, 14 mg/L/d and 1.2, respectively. Similarly, the individual EAA levels rose in NL6 from start to end (Fig 2C). The photo-acclimation response of microalgae towards excess light has been well-described, and upregulation of EAA during photo-acclimation was found in both microalgae and plants in response to high light exposure (Davis et al., 2013; Galili et al., 2016). Davis et al., (2013) suggest that the elevation of EAA could be due to an induction of *de novo* amino acid biosynthesis and/or protein catabolism from photodamage. Nevertheless, the fact that EAA content in NL4 (start and end) was not elevated might be related to their extreme N deficiency (Table 2). As stated, when N deprivation and high light were both applied to *Dunaliella tertioecta*, most amino acids were significantly decreased (Lee et al., 2014).

3.5 Optimized biomass with both high-quality protein and carotenoids

As the parameters indicating protein quantity, protein quality and carotenoid content varied substantially depending on the cultivating conditions, Fig 3 and Table 4 elucidate the effects of N and light intensity on the biomass quality of *D. salina*. Nitrogen concentration in the medium and light intensity, but particularly their interaction, significantly affected EAAI (Table 4; Fig 3B, 3C). These findings suggest that short N starvation and higher light intensity together enhance EAA production in *D. salina*, rather than either factor alone. Similar results were estimated for carotenoids content, where highest content was found with a combination of decreasing N and increasing light intensity, while only N starvation and not higher light showed also a significant separate effect (Table 4). It is noted however, that the correlation of light intensity to carotenoids

content was limited by the low sample sizes. When the effect of light intensity on carotenoid contents was analyzed from more samples obtained from this study, it indeed becomes evident that light intensity is correlated to carotenoid contents (Table 4; Fig 3G). Together, these results indicate that short N starvation and subsequent higher light intensity together are favored for carotenoids accumulation.

Overall, short N starvation together with higher light intensity are beneficial for the production of both EAA and carotenoids. Specifically, treatment NL4 (end) presented exceptional EAA enhancement and carotenoids accumulation in *D. salina*, indicating the possibility to produce microalgal biomass with both high protein quality and anti-oxidant strength. Consequently, *D. salina* demonstrated to be a valuable, and potentially unique species as novel protein source with high anti-oxidant activity for humans. Having been successfully applied in commercial production of *D. salina* using two-phase cultivation system, minor modifications can be applied to optimize biomass value, where short N starvation should be reached in the stationary phase in phase-one and a higher-light treatment should be applied in phase-two.

Broadly looking at the market of microalgal products, microalgae are mainly supplied as food supplement with high protein content and other nutritional values like vitamins.

Specifically for *D. salina*, the global market is mainly for high-value food and feed supplement. To the authors' knowledge, there is no microalgal product in the food industry which provides both high-quality protein and carotenoids at the same time. The results from this study thus present a step further towards production of microalgae with

multiple high-value compounds, which largely improves the production efficiency. The biomass can therefore be a superior source for food supplement and food ingredient, either consumed next to food sources or be incorporated into food products.

Conclusions

Protein quality and carotenoids content of *D. salina* could be simultaneously enhanced applying a two-phase cultivation strategy. Short N starvation should be reached in the stationary phase in phase-one to upregulate EAA production, and phase-two should be given higher illumination to boost carotenoids production. The optimized cultivation conditions resulted in production of *D. salina* biomass with an EAAI of 1.1 and cellular carotenoids content of 24 pg/cell. This study reveals that *D. salina* is a valuable, and potentially unique species bringing together both high-quality protein and carotenoid, thus can be used as protein source with antioxidant pro-vitamin A effect for humans.

Acknowledgements

This work was supported by the China Scholarship Council (File No. 201507650015) and the MIP i-Cleantech Flanders (Milieu-innovatieplatform; Environment innovation platform) project Microbial Nutrients on Demand (MicroNOD). Nico Helmsing from NIOO-KNAW is acknowledged for technical support.

Reference

Aasen, A.J., Eimhjellen, K.E., Liaaen-Jensen, S., Heinegård, D., Balaban, A.T., Craig, J.C., 1969. An Extreme Source of beta-Carotene. *Acta Chem. Scand.* 23, 2544–2545.

- Agusti, S., Kalff, J., 1989. The influence of growth conditions on the size dependence of maximal algal density and biomass. *Limnol. Oceanogr.* 34, 1104–1108.
- Alipanah, L., Rohloff, J., Winge, P., Bones, A.M., Brembu, T., 2015. Whole-cell response to nitrogen deprivation in the diatom *Phaeodactylum tricornutum*. *J. Exp. Bot.* 66, 6281–6296.
- Angell, A.R., Mata, L., de Nys, R., Paul, N.A., 2014. Variation in amino acid content and its relationship to nitrogen content and growth rate in *Ulva ohnoi* (Chlorophyta). *J. Phycol.* 50, 216–226.
- APHA, 2012. Standard Methods for Examination of Water and Wastewater. Am. Public Heal. Assoc. Washington, DC, USA.
- Becker, E.W., 2007. Micro-algae as a source of protein. *Biotechnol. Adv.* 25, 207–210.
- Ben-Amotz, A., 1995. New mode of *Dunaliella* biotechnology: two-phase growth for β -carotene production. *J. Appl. Phycol.* 7, 65–68.
- Ben-Amotz, A., 1993. Production of β -Carotene and Vitamins by the Halotolerant Alga *Dunaliella*. *Mar. Biotechnol.* 1, 411–417.
- Boland, M.J., Rae, A.N., Vereijken, J.M., Meuwissen, M.P.M., Fischer, A.R.H., van Boekel, M.A.J.S., Rutherford, S.M., Gruppen, H., Moughan, P.J., Hendriks, W.H., 2013. The future supply of animal-derived protein for human consumption. *Trends Food Sci. Technol.* 29, 62–73.

- Borowitzka, M.A., 1988. Algal growth media and sources of cultures, in: Micro-Algal Biotechnology. Cambridge University Press, Cambridge, pp. 456–465.
- Brown, N., Shilton, A., 2014. Luxury uptake of phosphorus by microalgae in waste stabilisation ponds: Current understanding and future direction. *Rev. Environ. Sci. Biotechnol.*
- Chen, C.Y., Lee, P.J., Tan, C.H., Lo, Y.C., Huang, C.C., Show, P.L., Lin, C.H., Chang, J.S., 2015. Improving protein production of indigenous microalga *Chlorella vulgaris* FSP-E by photobioreactor design and cultivation strategies. *Biotechnol. J.* 10, 905–914.
- Cole, A.J., Angell, A.R., de Nys, R., Paul, N.A., 2015. Cyclical changes in biomass productivity and amino acid content of freshwater macroalgae following nitrogen manipulation. *Algal Res.* 12, 477–486.
- Corcoran, E., Nellesmann, C., Baker, E., Bos, R., Osborn, D., Savelli, H., 2010. Wastewater and global change, in: Sick Water? The Central Role of Wastewater Management in Sustainable Development. United Nations Environment Programme, UN-HABITAT, GRID-Arendal, pp. 49–51.
- Davis, M.C., Fiehn, O., Durnford, D.G., 2013. Metabolic acclimation to excess light intensity in *Chlamydomonas reinhardtii*. *Plant, Cell Environ.* 36, 1391–1405.
- Dortch, Q., Clayton, J.R., Thoresen, S.S., Ahmed, S.I., 1984. Species differences in accumulation of nitrogen pools in phytoplankton. *Mar. Biol.* 81, 237–250.

- Dortch, Q., Clayton, J.R., Thoreson, S.S., Bressler, S.L., Ahmed, S.I., 1982. Response of marine phytoplankton to nitrogen deficiency: Decreased nitrate uptake vs enhanced ammonium uptake. *Mar. Biol.* 70, 13–19.
- Fabregas, J., Abalde, J., Cabezas, B., Herrero, C., 1989. Changes in protein, carbohydrates and gross energy in the marine microalga *Dunaliella tertiolecta* (Butcher) by nitrogen concentrations as nitrate, nitrite and urea. *Aquac. Eng.* 8, 223–239.
- Galili, G., Amir, R., Fernie, A.R., 2016. The Regulation of Essential Amino Acid Synthesis and Accumulation in Plants. *Annu. Rev. Plant Biol.* 67, 153–178.
- Gao, B., Liu, J., Zhang, C., Van de Waal, D.B., 2018. Biological stoichiometry of oleaginous microalgal lipid synthesis: The role of N:P supply ratios and growth rate on microalgal elemental and biochemical composition. *Algal Res.* 32, 353–361.
- Guerra, L.T., Levitan, O., Frada, M.J., Sun, J.S., Falkowski, P.G., Dismukes, G.C., 2013. Regulatory branch points affecting protein and lipid biosynthesis in the diatom *phaeodactylum tricornutum*. *Biomass and Bioenergy* 59, 306–315.
- Halsey, K.H., Milligan, A.J., Behrenfeld, M.J., 2011. Linking time-dependent carbon-fixation efficiencies in *Dunaliella Tertiolecta* (Chlorophyceae) to underlying metabolic pathways. *J. Phycol.* 47, 66–76.
- Kent, M., Welladsen, H.M., Mangott, A., Li, Y., 2015. Nutritional evaluation of Australian microalgae as potential human health supplements. *PLoS One* 10, e0118985.

- Kiran, B., Pathak, K., Kumar, R., Deshmukh, D., Rani, N., 2016. Influence of varying nitrogen levels on lipid accumulation in *Chlorella* sp. *Int. J. Environ. Sci. Technol.* 13, 1823–1832.
- Kiyota, H., Hirai, M., Ikeuchi, M., 2014. NblA1/A2-Dependent Homeostasis of Amino Acid Pools during Nitrogen Starvation in *Synechocystis* sp. PCC 6803. *Metabolites* 4, 517–531.
- Lamers, P.P., Janssen, M., De Vos, R.C.H., Bino, R.J., Wijffels, R.H., 2012. Carotenoid and fatty acid metabolism in nitrogen-starved *Dunaliella salina*, a unicellular green microalga. *J. Biotechnol.* 162, 21–27.
- Lamers, P.P., Janssen, M., De Vos, R.C.H., Bino, R.J., Wijffels, R.H., 2008. Exploring and exploiting carotenoid accumulation in *Dunaliella salina* for cell-factory applications. *Trends Biotechnol.* 26, 631–638.
- Lamers, P.P., Van De Laak, C.C.W., Kaasenbrood, P.S., Lorier, J., Janssen, M., De Vos, R.C.H., Bino, R.J., Wijffels, R.H., 2010. Carotenoid and fatty acid metabolism in light-stressed *Dunaliella salina*. *Biotechnol. Bioeng.* 106, 638–648.
- Lee, S.Y., Kim, S.H., Hyun, S.H., Suh, H.W., Hong, S.J., Cho, B.K., Lee, C.G., Lee, H., Choi, H.K., 2014. Fatty acids and global metabolites profiling of *Dunaliella tertiolecta* by shifting culture conditions to nitrate deficiency and high light at different growth phases. *Process Biochem.* 49, 996–1004.
- Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: pigments of photosynthetic

biomembranes. *Methods Enzymol.* 148, 350–382.

Lv, H., Cui, X., Tan, Z., Jia, S., 2017. Analysis of metabolic responses of *Dunaliella salina* to phosphorus deprivation. *J. Appl. Phycol.* 29, 1251–1260.

Marín, N., Morales, F., Lodeiros, C., Tamigneaux, E., 1998. Effect of nitrate concentration on growth and pigment synthesis of *Dunaliella salina* cultivated under low illumination and preadapted to different salinities. *J. Appl. Phycol.* 10, 405–411.

Markou, G., Chatzipavlidis, I., Georgakakis, D., 2012. Effects of phosphorus concentration and light intensity on the biomass composition of *Arthrospira (Spirulina) platensis*. *World J. Microbiol. Biotechnol.* 28, 2661–2670.

Markwell, M.A.K., Haas, S.M., Bieber, L.L., Tolbert, N.E., 1978. A modification of the Lowry procedure to simplify protein determination in membrane and lipoprotein samples. *Anal. Biochem.* 87, 206–210.

Muys, M., Sui, Y., Schwaiger, B., Lesueur, C., Vandenneuvel, D., Vermeir, P., Vlaeminck, S.E., 2019. High variability in nutritional value and safety of commercially available *Chlorella* and *Spirulina* biomass indicates the need for smart production strategies. *Bioresour. Technol.* 275, 247–257.

Obata, T., Fernie, A.R., 2012. The use of metabolomics to dissect plant responses to abiotic stresses. *Cell. Mol. Life Sci.* 69, 3225–3243.

Ogbonna, J.C., Tanaka, H., 1996. Night biomass loss and changes in biochemical

composition of cells during light/dark cyclic culture of *Chlorella pyrenoidosa*. J.

Ferment. Bioeng. 82, 558–564.

Oser, B.L., 1959. An Integrated Essential Amino Acid Index for Predicting the Biological Value of Proteins, Protein and Amino Acid Nutrition. Academic Press, Inc.

Phenomenex, 2003. Phenomenex EZ:faast amino acid analysis. Phenomenex, 411 Madrid Avenue, Torrance, CA 90501–1430, USA.

Raja, R., Hemaiswarya, S., Rengasamy, R., 2007. Exploitation of *Dunaliella* for β -carotene production. Appl. Microbiol. Biotechnol. 74, 517–523.

Remmers, I.M., D'Adamo, S., Martens, D.E., de Vos, R.C.H., Mumm, R., America, A.H.P., Cordewener, J.H.G., Bakker, L. V., Peters, S.A., Wijffels, R.H., Lamers, P.P., 2018. Orchestration of transcriptome, proteome and metabolome in the diatom *Phaeodactylum tricornutum* during nitrogen limitation. Algal Res. 35, 33–49.

Sanz-Luque, E., Chamizo-Ampudia, A., Llamas, A., Galvan, A., Fernandez, E., 2015. Understanding nitrate assimilation and its regulation in microalgae. Front. Plant Sci. 6, 899.

Sommer, A., 2001. Vitamin A Deficiency. Encycl. Life Sci.

Sui, Y., Muys, M., Vermeir, P., D'Adamo, S., Vlaeminck, S.E., 2019. Light regime and growth phase affect the microalgal production of protein quantity and quality with *Dunaliella salina*. Bioresour. Technol. 275, 145–152.

- Sui, Y., Vlaeminck, S.E., 2019. Effects of salinity, pH and growth phase on the protein productivity by *Dunaliella salina*. J. Chem. Technol. Biotechnol. 94, 1032–1040.
- Van de Waal, D.B., Ferrerueta, G., Van Donk, E., Huisman, J., Visser, P.M., Tonk, L., Matthijs, H.C.P., 2010. Pulsed nitrogen supply induces dynamic changes in the amino acid composition and microcystin production of the harmful cyanobacterium *Planktothrix agardhii*. FEMS Microbiol. Ecol. 74, 430–438.
- WHO/FAO/UNU Expert Consultation, 2007. Protein and amino acid requirements in human nutrition. World Health Organ. Tech. Rep. Ser. 1–265.
- Xie, T., Xia, Y., Zeng, Y., Li, X., Zhang, Y., 2017. Nitrate concentration-shift cultivation to enhance protein content of heterotrophic microalga *Chlorella vulgaris*: Over-compensation strategy. Bioresour. Technol. 233, 247–255.
- Zachleder, V., Bišová, K., Vítová, M., 2016. The Cell Cycle of Microalgae, in: Borowitzka, M.A., Beardall, J., Raven, J.A. (Eds.), The Physiology of Microalgae. Springer International Publishing, Cham, pp. 3–46.
- Zhang, Y., Liu, Y., Cao, X., Gao, P., Liu, X., Wang, X., Zhang, J., Zhou, J., Xue, S., Xu, G., Tian, J., 2016. Free amino acids and small molecular acids profiling of marine microalga *Isochrysis zhangjiangensis* under nitrogen deficiency. Algal Res. 13, 207–217.
- Zhuang, L.L., Azimi, Y., Yu, D., Wu, Y.H., Hu, H.Y., 2018. Effects of nitrogen and phosphorus concentrations on the growth of microalgae *Scenedesmus*. LX1 in suspended-solid phase photobioreactors (ssPBR). Biomass and Bioenergy 109, 47–53.

Figure captions

Fig. 1. Effect of two-phase cultivation on *D. salina*: (A) cell number and cell volume, (B) suspension protein and cellular protein, (C) suspension carotenoids and cellular carotenoids, (D) protein productivity and (E) carotenoids productivity. Different N levels were applied in phase-one and both N addition and higher illumination were applied in phase-two. Cultivation occurred at 25°C and pH 7.8-8. Data are expressed as means \pm standard deviation (n = 3)

Fig. 2. Effect of two-phase cultivation on *D. salina* at the end of each phase: (A) EAAI, EAA content and EAA productivity, (B) individual EAA level in treatment N4, NN4 (start and end) and NL4 (start and end) and (C) individual EAA level in treatment N6, NN6 (start and end) and NL6 (start and end). Different N levels were applied in phase-one and both N addition and higher illumination were applied in phase-two. Cultivation occurred at 25°C and pH 7.8-8. Data are expressed as means \pm standard deviation (n = 3)

Fig. 3. N and light intensity on EAAI and carotenoids content in ten *D. salina* samples: (A) N and light intensity vs. EAAI, (B) N vs. EAAI, (C) light intensity vs. EAAI, (D) N and light intensity vs. carotenoids content, (E) N vs. carotenoids content and (F) light intensity vs. carotenoids content. Highlighted blue dots represent all conditions where an EAAI above 1 was obtained. The corresponding carotenoids content obtained from above conditions were highlighted as orange dots. Light intensity vs. carotenoids content from 24 *D. salina* samples was presented in (G).

Fig. 1.




Fig. 2.

ACC






Fig. 3.



ACCEPTED



ACCEPTED MANUSCRIPT

Highlights

- *D. salina* can co-produce high quality protein and β -carotene for human nutrition
- Short nitrogen starvation promotes essential amino acid (EAA) production
- The essential amino acid index (EAAI) of *D. salina* can surpass FAO/WHO reference
- High light intensity can enhance β -carotene production
- A defined two-phase cultivation is beneficial for an optimized biomass composition of *D. salina*

ACCEPTED MANUSCRIPT